


Contents available at ISC and SID

Journal homepage: [www.rangeland.ir](http://www.rangeland.ir)



---

**Research and Full Length Article:**

## **Evaluation of Soil Quality Based on Minimum Data Set in Karvan Rangeland, Isfahan, Iran**

Habib Yazdanshenas<sup>A</sup>, Mohammad Jafari<sup>B</sup>, Hossein Azarnivand<sup>C</sup>, Hossein Arzani<sup>D</sup>

<sup>A</sup>M.Sc. Student of Range Management, Faculty of Natural Resources, University of Tehran, Iran

<sup>B</sup>Professor, Faculty of Natural Resources, University of Tehran, Iran (Corresponding Author),  
Email: jafary@ut.ac.ir

<sup>C,D</sup>Professor, Faculty of Natural Resources, University of Tehran, Iran

Received on: 20/02/2015

Accepted on: 29/05/2015

**Abstract.** Plant life and food production for human closely depend on fertile and healthy soil. Knowledge about qualitative properties of soil and its potential production can contribute us in the plantation, fertilization, utilization and land management. In addition, Rangelands cover a very large portion of the earth's surface and play an important role in food security and other ecosystem services. Therefore, the present study has been conducted in order to evaluate soil quality according to minimum data set in Karvan rangeland which is located in the west of Isfahan province, Iran. For this aim, three vegetation types including *Scariola orientalis-Astragalus gossypinus* (*Sc.or-As.go*), *Psathyrostachys fragile-Astragalus gossypinus* (*Psa.fr-As.go*) and *Cousinia bachtiarica-Astragalus gossypinus* (*Cu.ba-As.go*) were selected in the study area. Then, four transects were established by a random systematic sampling; bias was placed to the general and lateral slope at the each vegetation type. The soil samples were taken at the start and end of each transect from two different depths (0-20 and 20-75 cm). Soil samples were analyzed and the physicochemical factors such as texture (silt, clay and sand), pH, Nitrogen (N), Phosphorus (P), potassium (K), Organic Carbon (OC) and Organic Matter (OM) were measured. Then, Soil Quality (SQ) indices were calculated using Bajracharya formula. The results showed that soil of the study area had a poor quality and also, there was different soil qualities regarding three vegetation types. It was found that OC (or OM) and N had maximum limitations on soil of Karvan region and caused low SQ indices. Also, pH only without any limitations was put in the highest rank for SQ measuring.

**Key words:** Soil quality, Minimum data set, Vegetation and fertility, Karvan region

## Introduction

Soil is one of the most important natural elements that ensure plant growth and provide more than 97% of the world's food requirements (Agheli Kohan and Sadeghi, 2004). Also, it plays an important role as a seed bed and provides all the requirements for plants (Raymond *et al.*, 1998). For example, the crop harvest was followed by the erosion and depletion in soil elements. This has led to the reduction of some important elements such as nitrogen (70%), potassium (90%) and phosphorous (100%) (Dreshcel *et al.*, 2001). Therefore, Soil management could change the dynamics of soil matter and seasonal and spatial distributions by changing the quality and quantity of plant residues, and the nutrient element in the soil (Kandeler *et al.*, 1999). Physical and chemical properties of soil and fertility are used to investigate the sustainable land potential (Lynn *et al.*, 2009). There is a difference concerning soil properties and elements (except P) between rangeland and cultivated lands (Zehtabian *et al.*, 2004) and some of the soil characteristics will be destroyed due to the land use changes for agricultural purposes (Ahmadi Ilakhchi *et al.*, 2002). For example, soil organic carbon reservoir has been influenced severity by changing the land use and land management (Lorenz *et al.*, 2008) or storage capacity of soil has been effectively related to some factors such as clay, silt and sand contents and porosity index (Rezaei, 2003) so that all soil properties affect soil productivity. However, rangelands have an important role in economy of society and culture (Rezaei *et al.*, 2006a). Thus, high quality range plant production needs appropriate soil and enough elements which are available for plants and there should be a balance among the elements in soil (Tandon, 1989). There are other methods to evaluate the ecosystem and soil such as landscape functional analysis (LFA) which needs some simple factors

including soil cover, litter cover, cryptogam cover, shell brittle, erosion properties, sediment, micro-topography, quenched test and soil texture (Tongway and Hindley, 2004).

In all methods, adequate information may be necessary for quality assessments (Andrews *et al.*, 2002). Soil quality indexing is a new approach in spatial and temporal evaluation of land management system effects on soil capacity to function (Erkossa *et al.*, 2007). Soil Quality Indices (SQI) are regarded as a complex set of physicochemical soil factors; they are easily calculated and have the following features: being sensitive to land management changes, being simply measured, having continuity along the site and at all time, closely investigating and measuring, very cheap and adaptable for all ecosystems (Schoenholtza *et al.*, 2000). Soil quality (SQ) affects the rangeland (USDA, 2001) through plant production, reproduction, mortality, erosion, water production and water quality, wildlife habitat, carbon sequestration, vegetation changes, establishment and growth of invasive plants and rangeland health. Soil quality can be also defined for such issues as productivity, environmental quality or human health. SQ is a concept that is hidden amidst the statements of all soil scientists (SBSMNFR, 2008) and includes the assessment of soil properties and ability while surveying the relationship between their processes as a part of healthy ecosystem (Schoenholtza *et al.*, 2000). Nowadays, most researchers only use physical or chemical properties to evaluate the SQ; however, considering that most of soil potentials are related to chemical, physical and biological factors, a complex of all factors should be used to calculate the SQI (Barrios *et al.*, 2006). Rangeland re-vegetation success not only depends on climate, topography and management, but is strongly influenced by soil (Heady and Child, 1994). There are many different parameters and



**Table 1.** Soil parameters and ranking values for evaluating them (NRC, 1993)

Parameters	Ranking Values				
	0.2	0.4	0.6	0.8	1
Soil pH	<4	4.1-4.9	5-5.9	6-6.4	6.5-7.5
Soil organic carbon (%)	<0.5	0.6-1	1.1-2	2.1-4	>4
Fertility (NPK)	Low	Mod Low	Moderate	Mod. High	High
Soil textural class	C, S	CL, SC, SiC	Si, LS	L, SiL, SL	SiCL, SC
SQI	V. poor	Poor	Fair	Good	Best

Abbreviation= SQI: Soil Quality Index, C:Clay, S:Sand, CL: Clay loam, SC: Sandy Clay, SiC: Silty Clay, Si: Silt, LS: Loamy sand, SiL: Silty loam, SL: Sandy loam, LS: Loamy Sand, SL: Sandy loam, SiCL: Silty clay loam and SCL: Sandy Clay loam

SQI was measured for the first and second depths of soil separately. Then, SQI for total soil profile (75 cm) was measured according to top and subsoil QIs as follows (Equation 2):

$$SQI = [(20 \times SQI) + (55 \times SQI)] / 75 \quad \text{(Equation 2)}$$

Also, in order to determinate the difference between soil physicochemical properties and SQI in two different depths and among three vegetation types, the Kolmogorov-Smirnov test (KS-test) was used to identify the data normality and then, the One-way ANOVA test and the post hoc test were performed using the Duncan method for grouping treats by the means of SPSS software (17.0 version).

**Results**

The initial comparison of soil physicochemical properties of three

vegetation types showed that there were significant differences ( $p \leq 0.01$ ) between some factors such as OC, P, K and Clay percent that could affect SQ. Regarding gravel, sand and silt percent, there were no significant differences among three vegetation types at two depths of soil profile. But for OM, the highest value was obtained in *Scariola orientalis-Astragalus gossypinus* type ( $p \leq 0.01$ ). ANOVA table shows that SQI was not significantly different ( $p \leq 0.01$ ) at different depths of soil in the study area except first depth at 5% confidence (Table 2). Table 3, shows means comparisons between factors based on Duncan test. Also Table 4 show soil factors, ranking values and soil quality index

**Table 2.** Analysis of variance for SQI at different soil depths among vegetation types

Depth	S.O.V	df	SS	MS	Sig.	F
(0-20)	Between Groups	2	0.163	0.082	0.001*	7.05*
	Within Groups	21	0.090	0.004	-	-
	Total	23	0.253	-	-	-
(20-75)	Between Groups	2	0.049	0.25	0.021 <sup>ns</sup>	4.6 <sup>ns</sup>
	Within Groups	21	0.092	0.004	-	-
	Total	23	0.142	-	-	-

S.O.V: Sources of variation, df: Degrees of freedom, SS: Sum of Squares, MS: Mean-Squares, \*: Significant at 1% confidence

**Table 3.** Mean values of important soil physicochemical properties and their difference at first and second depths of soil

Vegetation types	Depth	Gravel (%)	Sand (%)	Silt (%)	Clay (%)	OC (%)	OM (%)	pH	N (Meq/L)	P (Meq/L)	K (Meq/L)	SQI
<i>Scariola orientalis-Astragalus gossypinus</i>	d <sub>1</sub> (0-20)	32.70 <sup>ns</sup>	22.75 <sup>ns</sup>	32.90 <sup>ns</sup>	34.40 <sup>ns</sup>	0.56 <sup>a</sup>	0.93 <sup>ns</sup>	7.87 <sup>ns</sup>	4.40 <sup>ns</sup>	16.80 <sup>ab</sup>	76.40 <sup>a</sup>	0.54 <sup>a</sup>
<i>Psathyrostachys fragile-Astragalus gossypinus</i>		33.00 <sup>ns</sup>	27.20 <sup>ns</sup>	42.40 <sup>ns</sup>	30.40 <sup>ns</sup>	0.45 <sup>b</sup>	0.72 <sup>ns</sup>	7.90 <sup>ns</sup>	4.20 <sup>ns</sup>	15.50 <sup>b</sup>	61.50 <sup>b</sup>	0.42 <sup>b</sup>
<i>Cousinia bachtiarica-Astragalus gossypinus</i>		34.60 <sup>ns</sup>	34.60 <sup>ns</sup>	32.10 <sup>ns</sup>	34.40 <sup>ns</sup>	0.55 <sup>a</sup>	0.94 <sup>ns</sup>	7.89 <sup>ns</sup>	4.60 <sup>ns</sup>	19.50 <sup>a</sup>	80.20 <sup>a</sup>	0.43 <sup>b</sup>
<i>Scariola orientalis-Astragalus gossypinus</i>	d <sub>2</sub> (20-75)	38.50 <sup>ns</sup>	38.50 <sup>ns</sup>	33.10 <sup>ns</sup>	38.60 <sup>b</sup>	0.54 <sup>a</sup>	0.90 <sup>a</sup>	7.88 <sup>b</sup>	4.60 <sup>ns</sup>	13.80 <sup>ns</sup>	60.50 <sup>b</sup>	0.40 <sup>ns</sup>
<i>Psathyrostachys fragile-Astragalus gossypinus</i>		47.50 <sup>ns</sup>	30.90 <sup>ns</sup>	36.90 <sup>ns</sup>	32.10 <sup>a</sup>	0.42 <sup>b</sup>	0.68 <sup>b</sup>	7.89 <sup>b</sup>	4.10 <sup>ns</sup>	16.30 <sup>ns</sup>	57.50 <sup>b</sup>	0.32 <sup>ns</sup>
<i>Cousinia bachtiarica-Astragalus gossypinus</i>		42.20 <sup>ns</sup>	42.20 <sup>ns</sup>	31.60 <sup>ns</sup>	31.10 <sup>ab</sup>	0.25 <sup>b</sup>	0.43 <sup>c</sup>	7.97 <sup>a</sup>	4.00 <sup>ns</sup>	17.00 <sup>ns</sup>	71.60 <sup>a</sup>	0.41 <sup>ns</sup>

Similar letters had no significant difference at first and second soil layers among three vegetation types (p≤0.01)

**Table 4.** Soil factors, ranking values and Soil Quality Index (SQI)

Vegetation Types	Parameters	Ranking Values				
		0.2	0.4	0.6	0.8	1
<i>Scariola orientalis-Astragalus gossypinus</i>	Soil pH	-	-	-	-	7.87
	Soil organic carbon(%)	-	0.55	-	-	-
	Fertility (NPK)	-	Mod Low	-	-	-
	Soil textural class	-	Clay loam	-	-	-
	SQI	-	-	<b>Fair</b>	-	-
<i>Psathyrostachys fragile-Astragalus gossypinus</i>	Soil pH	-	-	-	-	7.90
	Soil organic carbon(%)	0.41	-	-	-	-
	Fertility (NPK)	low	-	-	-	-
	Soil textural class	-	-	-	-	Silty clay loam
	SQI	-	<b>Poor</b>	-	-	-
<i>Cousinia bachtiarica-Astragalus gossypinus</i>	Soil pH	-	-	-	-	7.93
	Soil organic carbon (%)	0.40	-	-	-	-
	Fertility (NPK)	low	-	-	-	-
	Soil textural class	-	Clay loam	-	-	-
	SQI	-	<b>Poor</b>	-	-	-

Investigation of soil physicochemical properties in this region indicated that there is no restriction about some factors. For example, pH value belongs to high value ranking in all three vegetation types in the study area.

In general, soil properties in *Psa. fr-As. go* and *Co. ba-As. go* types had similarities that were much closer together in some cases such as fertility. Soil of *Sc. or-As. go* type was more nutritious than two other vegetation types. Soil texture in three vegetation types was mostly put in heavy texture ranking. For SQI, only *Scariola orientalis-Astragalus gossypinus* had fair ranking among three vegetation types and had shown a better condition (Table 3).

## Discussion

Research and adequate information about SQ could contribute us in managing the land uses. Soil testing is the most accurate method for determining whether sufficient nutrient is present. SQ is the capacity of a specific kind of soil to function within natural or managed ecosystem boundaries, sustain plant and animal productivity, maintain or enhance the quality of water and air and support human health and habitation (USDA, 2001). Numerous factors could affect SQ. Some proceedings such as range management can change the SQ although Steffens *et al.* (2011) reported that some factors including livestock grazing had no effects on the amount of OC (OM and SQ) in their study site. However, SQ parameters will be changed even without human interventions (SBSMNFR, 2008). Climate factor could affect SQ (Vallejo *et al.*, 2005) and will be most likely the most important factor. Also, some lateral factors could have less effect on SQ. Rezaee *et al.* (2006) found that geographic aspect had no effects on soil properties in the study area. The result of this study showed that there were moderate-low and low ranking values of soil fertility and then poor and fair SQ

(Tables 4&3). This can be different not only around the world but also in different vegetation types (similar to this research). Present study indicates that SQI related to soil properties was not in good conditions among all vegetation types in which SQI was between 0.32 to 0.54 (Table 3). Awasthi (2004) reported that SQI had the highest and lowest values (0.69 and 0.17) in the undisturbed forest lands and Khet land in Mardi watershed of Middle Mountain, respectively. SQ specifies that there is a kind of fertilizer and tillage required in soil. But, it is noteworthy that the nutrient elements in plant and soil structure vary at different times of the year (Demirosoy *et al.*, 2010). Therefore, SQI should be measured and used over time based on our ability and perception (Dumanski and Pieri, 2000).

Jagadamma and Lal (2010) reported that there is OM available in heavy soil texture more than light texture; these factors can influence soil quality that is different with the present study. First vegetation type (*Scariola orientalis-Astragalus gossypinus*) had a lower clay percent but the highest OM percent. Also, He *et al.* (2009) indicated that the amount of OC was higher in clay texture in different soil depths. In present research, soil had a heavy texture in three vegetation types and soil components were relatively in equal Ratio. Other researchers reported that physical soil factors had more effects than chemical properties on soil potential, dry matter production and overall forage yield for animals (Rezaei *et al.*, 2006b). Zornoza *et al.* (2007) concluded that Organic Carbon (OC) had the most effects on SQ in the study area in Alicante province, Spain. However, in present study, physicochemical soil factors have been combined while influencing SQI. Also, because of the region location which was located in boundary of steppe and semi-steppe zone, some soil properties such as OC, OM and N percent were very low

(Table 3). SQI is increasingly proposed as an integrative indicator of environmental quality (NRC, 1993); therefore, soil status in this region is not in a good condition. But soil pH was good among three vegetation types in the study area. In general, the main reason for low SQ in the study area returns to climate. Due to location in the arid and semi-arid land, Karvan rangeland has low annual precipitation. As a result, rainfall and humidity affect plant vegetation and vegetation in turn influences soil physicochemical properties. SQI among three vegetation types had no significant difference at 1% confidence but *Sc.or-As.go* type had the highest SQI (0.52) as compared to the other vegetation types ( $p \leq 0.05$ ). Similar to our result, Bajracharya et al. (2006) reported that SQI land uses were not distinctly different in a mid-hill watershed of Nepal.

### Conclusion

The results of present study showed that soil quality was placed in the poor class based on minimum data set method using Bajracharya formula and NRC ranking. Chemical properties of soil such as OC and N content were the main reasons for low quality in this region although pH was in the best ranking. Soil texture approximately imposed no limitations on soil quality. Also, SQI in *Sc. or-As. go* type was better than two other vegetation types in the study area.

### Acknowledgment

We would like to appreciate the local people of Karvan area and also special thanks to Mr. Tarnian, Mr. Yeganeh, Mr. Moameri, Mr. Nazarzadeh and Mr. Assadollahi for their help.

### Literature Cited

Agheli Kohan, L. and Sadeghi, H. 2004. Estimate the economic impact of soil erosion in Iran. *Jour. Economic Studies*, 15: 1-14.

Ahmadi Ilakhchi, A., Hajabbasi, M. A., and Jalalian, A., 2002. The effect of land use change

of rangeland to dry land farming on runoff production, lose and soil quality in Dorahan of Chaharmahal Bakhtiari province, *Jour. Sciences Techniques in Natural Resources*, 6(4): 103-115. (In Persian).

- Andrews, S. S., Karlen, D. L., and Mitchell, J. P., 2002. A comparison of soil quality indexing methods for vegetable production systems in Northern California. *Agriculture, Ecosystems & Environment*, 90(1): 25-45.
- Andrews, S. S. Carroll, C. R., 2001. Designing a decision tool for sustainable agroecosystem management: soil quality assessment of a poultry litter management case study. *Ecol. Applic.* 11(6): 1573-1585.
- Awasthi, K. D., 2004. Land use change effect on soil degradation, carbon and nutrient stocks and greenhouse gas emission in mountain watersheds. Agricultural university of Norway.
- Bajracharya, R. M., Sharma, S., Dahal, B. M., Sitaula, B. K., Rokaya, K., and Jeng, A., 2006. Assessment of soil quality using physiochemical and biological indicators in a mid-hill watershed of Nepal. In *Proceedings of Int'l. Seminar on Environmental and Social Impacts of Agricultural Intensification in Himalayan Watersheds* (pp. 105-114).
- Barrios, E., Delv, R. J., Bejunda, M., Mowo, J., Agunda, J., Ramisch, J., Trejo, M. T. and Thomas, R. J., 2006. Indicators of soil quality: a south-south development of a methodological guide for linking local and technical knowledge. *Geoderma*, 135: 248-259.
- Demirosoy, L., Demisosoy, H., Ersoy, B., Balci, G. and Kizlkaya, R., 2010. Seasonal Variation of N, P, K Ca content of leaf, crown and root of Sweet Charlie strawberry under different irrigation. *Zemdirbyste- Agriculture*, 97(1): 23-32.
- Dreshcel, P., Guiele, L., Kunze, D. and Cofie, O. 2001. Population Density, Soil Nutrient, Depletion and Economic Growth in Sub-Saharan Africa. *Ecological Economics*, 38: 251-258.
- Dumanski, J. and C. Pieri., 2000. Land quality indicators: research plan. *Agric. Ecosyst. Environ.* 81: 93-102.
- Erkossa, T., Itanna, F. and Stahr, K., 2007. Indexing soil quality: a new paradigm in soil science research. *Soil Research*, 45(2): 129-137.
- He, N., Wu, L., Wang, Y and Han, X., 2009. Changes in carbon and nitrogen in soil particle-size fractions along a grassland restoration chronosequence in northern China. *Geoderma*, 150: 302-308.

- Heady, F. H., and Child, D., 1994. *Rangeland Ecology and Management*. West view Press, San
- Jagadamma, S. and Lal, R., 2010. Distribution of organic carbon in physical fractions of soils as affected by agricultural management. *Biology and Fertility of Soils*, 46: 543-554.
- Kandeler, E., Stemmer, M. and Klimanek, E. M., 1999. Response of soil microbial biomass, urease and size fractions in no-till and meadow soils. *Biology and Fertility of Soils*, 44: 1043-1051.
- Lorenz, K., Lal, R. and Shipitalo, M. J., 2008. Chemical stabilization of organic carbon pools in particle xylanase within particle size fractions to long-term soil management. *Soil Biology and Biochemistry*, 31: 261-273.
- Lynn, I., Manderson, A., Page, M., Harmsworth, G., Eyles Douglas, G., Mackay, A. and Newsome, P. 2009. Land Use capability survive hand book. A New Zealand Handbook for the classification of land 3<sup>rd</sup> ed. Hamilton. Ag Research; Lincoln, Landcare research; Lower Hott, GNS Science. 163 p.
- NRC (National Research Council). 1993. Soil and water quality: an agenda for agriculture. National academy press, Washington, DC, 516 pp.
- Peterson, E. H., Pannel, D. G., Nordblom, T. L. and Shomo, F. 2002. Potential benefit from alternative areas of agricultural research for dry farming in northern Syria. *Jour. Agricultural System*. 11(2): 93-108.
- Raymod, W., Millen Roy, L. and Donahue, S. 1998. An Introduction to soil and plant growth, prentice-Hall. Sixth Edition, International (UK) limited, London.
- Rezaei, S. A., 2003. The use of a soil quality index in site capability assessment for extensive grazing. Ph.D Dissertation, University of Western Australia, Perth, Australia.
- Rezaei, S. A., Arzani, H. and Tongway, D., 2006a. Assessing rangeland capability in Iran using landscape function indices based on soil surface attributes. *Jour. Arid Environments*, 65: 460-473.
- Rezaei, S. A., Gilkes, R. J. and Andrews, S. S., 2006b. A minimum data set for assessing soil quality in rangelands. *Geoderma*, 136: 229-234.
- SBSMNFR (Scientific Background for Soil Monitoring on National Forests and Rangelands). 2008. Workshop Proceedings, April 29-30, Denver, Colorado.
- Schoenholtz, S. H., Van Miegroetb, H. and Burgerc, J.A., 2000. A review of chemical and physical properties as indicators of forest soil quality: challenges and opportunities. *Forest Ecology and Management*, 138: 335-356.
- Steffens, M., Kolbl, A., Schork, E., Gschrey, B. and Kogel Knabner, I., 2011. Distribution of soil organic matter between fractions and aggregate size classes in grazed semiarid steppe soil profiles. *Plant and Soil*, 338: 63-81.
- Tandon, H. L. S., 1989. Fertilizer management in food crop, Fertilizer development and consultation organization. *Jour. New Delhi India*, P. 191-198.
- Tongway, D. and Hindley, N. 2004. Landscape Function Analysis: Procedures for Monitoring and Assessing Landscapes CSIRO, Brisbane, Australia.
- USDA (The United States Department of Agriculture). 2001. Guidelines for soil quality assessment in conservation planning. Natural Resources Conservation Service (NRCS). Soil Quality Institute. 38 p.
- Vallejo, V. R., Diaz, Fierros, F. and Rosa, D., 2005. Impactos sobre los recursos edaficos. Evaluacion Preliminar de los Impactos en Espana por Efecto del Cambio Climatico. Ministerio de Medio Ambiente, pp. 355-398. (In Spanish).
- Zehtabian, Gh. R., Amiri, B. and Surrey, M., 2004. Comparison of soil Nutrient elements status, with emphasis on the NPK Provincial Bureau of Land and pasture (Case Study: Khodabandeh, Zanjan). Research and development in agriculture and horticulture. 68: 1-11. (In Persian).
- Zornoza, R., Mataix-Solera, J., Guerrero, C., Arcenegui, V., Mayoral, A. M., Morales, J. and Mataix-Beneyto, J., 2007. Soil properties under natural forest in the Alicante Province of Spain. *Geoderma*, 142: 334-341.

## ارزیابی کیفیت خاک در مراتع کرون (اصفهان، ایران) بر اساس مجموعه حداقل داده‌ها

حبیب یزدانشناس<sup>الف</sup>، محمد جعفری<sup>ب</sup>، حسین آذرینوند<sup>ج</sup>، حسین ارزانی<sup>د</sup>

<sup>الف</sup> دانشجوی کارشناسی ارشد مرتعداری، دانشکده منابع طبیعی، دانشگاه تهران

<sup>ب</sup> استاد گروه احیای مناطق خشک و کوهستانی، دانشکده منابع طبیعی، دانشگاه تهران (نگارنده مسئول)، پست الکترونیک: jafary@ut.ac.ir

<sup>ج</sup> <sup>د</sup> استاد گروه احیای مناطق خشک و کوهستانی، دانشکده منابع طبیعی، دانشگاه تهران

تاریخ دریافت: ۱۳۹۳/۱۲/۰۱

تاریخ پذیرش: ۱۳۹۴/۰۳/۰۸

**چکیده.** حیات نباتات و تولید غذا برای بشر ارتباط تنگاتنگ با سلامت و حاصلخیزی خاک دارد. دانش در خصوص خصوصیات کیفی خاک و پتانسیل آن می‌تواند برای کشت گیاهان، کوددهی، بهره‌برداری و مدیریت زمین مؤثر باشد. از سویی مراتع، سطح وسیعی از زمین را پوشانده‌اند و نقش بسزایی در امنیت غذایی و سایر خدمات اکوسیستم دارند. بدین ترتیب این مطالعه به منظور بررسی کیفیت فیزیکی‌شیمیایی خاک بر اساس حداقل داده در مراتع کرون در غرب استان اصفهان انجام شد. ابتدا سه تیپ گیاهی شامل *Psathyrostachys fragile-Astragalus* (Sc. or-As. go) *Scariola orientalis-Astragalus gossypinus* در منطقه انتخاب شدند. سپس چهار ترانسکت ۲۰۰ متری با فاصله ۱۰۰ متر از همدیگر و اریب با شیب عمومی و جانبی منطقه در هر تیپ گیاهی قرار داده شد. از ابتدا و انتهای هر ترانسکت و از دو عمق مختلف (۲۰-۷۵ و ۲۰-۷۵ سانتی‌متر) نمونه‌برداری انجام شد. نمونه‌های خاک مورد آزمایش قرار گرفت و خصوصیات بافت (شن، سیلت و رس)، اسیدیته، نیتروژن، فسفر، پتاسیم، کربن آلی و ماده آلی اندازه‌گیری شد. شاخص کیفیت خاک با استفاده از فرمول Bajracharya محاسبه شد. نتایج نشان داد که شاخص کیفیت خاک در این منطقه در وضعیت فقیر قرار دارد. همچنین این شاخص بین سه تیپ گیاهی منطقه متفاوت بود. خصوصیات مقادیر ماده آلی و ازت بیشترین محدود و تأثیر بر کاهش شاخص کیفیت خاک در منطقه نشان داده شد. همچنین تنها اسیدیته خاک بدون هیچگونه محدودیتی برای محاسبه شاخص کیفیت خاک در بالاترین سطح رده‌بندی قرار گرفت.

**کلمات کلیدی:** کیفیت خاک، مجموعه حداقل داده، پوشش و حاصلخیزی، ناحیه کرون